

s- and d-wave solution of Eliashberg equations with finite bandwidth

G.A. Ummarino and R.S. Gonnelli

INFN - Dipartimento di Fisica, Politecnico di Torino, c.so Duca degli Abruzzi 24, 10129 Torino, Italy

In this work, we discuss the results of the direct solution of the Eliashberg equations with finite bandwidth, in the cases of *s*- and *d*-wave symmetry for the pair wave function and in the presence of scattering from impurities. We show that the reduction of the critical temperature T_c due to the finite bandwidth depends on the value of the bandwidth itself, but is almost independent of the symmetry of the order parameter. The same happens for the shape of $Z(\omega)$ and $\Delta(\omega)$. Moreover, we discuss the effect of the finite bandwidth on the shape of the quasiparticle density of states. The results clearly indicate that the infinite bandwidth approximation leads to an underestimation of the electron-boson coupling constant.

High- T_c cuprates and fullerenes are characterized by phononic energies (Ω_{phon}) comparable with the electronic ones (E_F) while in low- T_c superconductors it is always $E_F \gg \Omega_{\text{phon}}$. This last condition leads to the standard Eliashberg equations [1] obtained within the limit $E_F \rightarrow +\infty$. In this paper, we study the effect of a finite bandwidth [2,3] on some relevant physical quantities in the framework of the Eliashberg theory. For simplicity, we disregard the important related problem of the breakdown of the Migdal's theorem, and put to zero the Coulomb pseudopotential μ^* [4]. The kernels of the Eliashberg equations (EE) for the renormalization function $Z(\omega, \mathbf{k})$ and the order parameter $\Delta(\omega, \mathbf{k})$ contain the retarded electron-boson interaction $\alpha^2(\Omega, \mathbf{k}, \mathbf{k}')F(\Omega)$. Referring to high- T_c cuprates, we assume $\mathbf{k}$ and $\mathbf{k}'$ to lie in the *ab* plane (CuO_2 plane) and call ϕ and ϕ' their azimuthal angles in this plane [3,5,6]. We integrate over the effective band normal to the Fermi surface from $-W$ to $+W$, and expand $\alpha^2(\Omega, \phi, \phi')F(\Omega)$ in terms of basis functions $\psi_0(\phi) = 1$ and $\psi_1(\phi) = \sqrt{2}\cos(2\phi)$ in the following way:

$$\alpha^2(\Omega, \phi, \phi')F(\Omega) = \alpha_{00}^2 F(\Omega)\psi_0(\phi)\psi_0(\phi') + \alpha_{11}^2 F(\Omega)\psi_1(\phi)\psi_1(\phi').$$

We then search for a *s* + *id* solution with

$$\Delta_n(\phi) \equiv \Delta(i\omega_n, \phi) = \Delta_s(i\omega_n) + \Delta_d(i\omega_n)\psi_1(\phi) \\ Z_n(\phi) \equiv Z(i\omega_n, \phi) = Z_s(i\omega_n) + Z_d(i\omega_n)\psi_1(\phi).$$

In this case, and in the Matsubara representation, the Eliashberg equations in the presence of

impurities become [2,3]:

$$\omega_n Z_n(\phi) = \omega_n + \frac{k_B T}{\pi} \sum_{m=-\infty}^{+\infty} \int_0^{2\pi} d\phi' \left[\lambda_{m,n}(\phi, \phi') N_m(\phi') \theta_m(\phi') + \frac{N_n(\phi') \theta_n(\phi')}{k_B T \tau} \right]$$

$$\Delta_n(\phi) Z_n(\phi) = \frac{k_B T}{\pi} \sum_{m=-\infty}^{+\infty} \int_0^{2\pi} d\phi' \left[\lambda_{m,n}(\phi, \phi') P_m(\phi') \theta_m(\phi') + \frac{P_n(\phi') \theta_n(\phi')}{k_B T \tau} \right]$$

$$\theta_n(\phi) = \tan^{-1} \left[W/2 \sqrt{\omega_n^2 Z_n^2(\phi) + \Delta_n^2(\phi) Z_n^2(\phi)} \right]$$

$$\lambda_{m,n}(\phi, \phi') = 2 \int_0^{+\infty} d\Omega \frac{\Omega \alpha^2(\Omega, \phi, \phi') F(\Omega)}{\Omega^2 + (\omega_n - \omega_m)^2}$$

where $\omega_n = \pi(2n+1)k_B T$, and τ^{-1} is the impurity scattering rate. Furthermore we know that:

$$P_n(\phi) = \Delta_n(\phi) Z_n(\phi) / \sqrt{\omega_n^2 Z_n^2(\phi) + \Delta_n^2(\phi) Z_n^2(\phi)}$$

$$N_n(\phi) = \omega_n Z_n(\phi) / \sqrt{\omega_n^2 Z_n^2(\phi) + \Delta_n^2(\phi) Z_n^2(\phi)}.$$

Thus we have four equations for $Z_s(i\omega_n)$, $Z_d(i\omega_n)$, $\Delta_s(i\omega_n)$ and $\Delta_d(i\omega_n)$. Here we only consider the case $Z_d(i\omega_n) \equiv 0$ [6].

In our numerical analysis we put, for simplicity, $\alpha_{11}^2 F(\Omega) = g_d \cdot \alpha_{00}^2 F(\Omega)$ where g_d is a constant and, as a consequence, $\lambda_d = g_d \cdot \lambda_s$ [3,6]. We used for $\alpha_{00}^2 F(\Omega)$ the spectral function we experimentally determined in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$ (BSCCO) [7]




Figure 1. (a) The critical temperature T_c versus the bandwidth W . (b) $\Delta(i\omega_{n=0})$, at $T=4$ K, versus W . The inset of (a) reports λ_{eff} versus W .

appropriately scaled to give, in both the s and d cases, $T_c = 97$ K in the $W = \infty$ limit. The coupling constants result to be $\lambda_s = 3.15$ for the s -wave case, $\lambda_s = 2$ and $\lambda_d = 2.3$ for the d -wave one. From the direct numerical solution of the EE we found that the symmetry of $\Delta_n(\phi)$ depends on the values of the coupling constants λ_s and λ_d and, for particular values of λ_s and λ_d , on the starting values of Δ_s and Δ_d .

As shown in Fig. 1 (a) and (b), the dependence of T_c and $\Delta(i\omega_{n=0})$ at $T=4$ K on the bandwidth W is almost the same in both the s - and d -wave symmetries. The reduction of the bandwidth results in a sensible reduction of T_c and, on a minor extent, of $\Delta(i\omega_{n=0})$. In Fig. 2 we show the d -wave normalized density of states, calculated through the analytical continuation with Padé approximants of the imaginary-axis solution, for various values of W at $T=2$ K [3]. As in the s -wave case




Figure 2. The d -wave normalized density of states, at $T = 2$ K, for $W=100, 350$ and 10000 meV.

(not shown), the effect of a small bandwidth is remarkable particularly at high energies. These curves are in *better agreement* with tunneling experimental data than those obtained for an infinite bandwidth, particularly for the presence of a *dip* above the peak (at about 2Δ) and because they asymptotically tend to 1 from above.

We have also calculated the real and imaginary parts of the functions $\Delta(\omega)$ and $Z(\omega)$ in the s - and d -wave cases. For small values of W , these functions are markedly modified with respect to the infinite bandwidth case, independently of the symmetry. In the inset of Fig. 1(a) we show (only in the s -wave case) the values of λ_{eff} necessary to obtain $T_c = 97$ K, for various values of W . We conclude that the use of the standard Eliashberg equations when $E_F \simeq \Omega_{phon}$ leads to *underestimate* the real value of the electron-boson coupling constant.

REFERENCES

1. G.M. Eliashberg, Sov. Phys. JETP 3 (1963) 696.
2. Han-Yong Choi, Phys. Rev. B 53 (1996) 8591.
3. G.A. Ummarino and R.S. Gonnelli, to be published.
4. G.A. Ummarino and R.S. Gonnelli, Phys. Rev. B 56 (1997) 14279.
5. C.T. Rieck, D. Fay and L. Tewordt, Phys. Rev. B 41 (1989) 7289.
6. G.A. Ummarino and R.S. Gonnelli, Physica C, in press.
7. R.S. Gonnelli *et al.*, Physica C 275 (1997) 162.